



SA-P 72.5%

GUARANTEED MINIMUM ANALYSIS

Soluble Potash (K ₂ O).....	27.5%
Salicylic Acid.....	72.5%

EZ-Gro SA-P 72.5% contains Salicylic Acid, a naturally occurring plant hormone, which plays a vital role in plant growth and development, photosynthesis, transpiration, ion uptake and transport. Salicylic Acid improves resistance to environmental stressors including drought, temperature, and salinity; and enhances flowering and fruit yield.

APPLICATION RATES

EZ-Gro SA-P 72.5% can be used as a foliar spray, a soil drench and through fertigation systems. It can be mixed with most fungicides, insecticides and other liquid fertilizers. Do not mix concentrates together. Add water to spray tank to ¾ full followed by the **EZ-Gro SA-P 72.5%**, pesticides and fertilizers. When mixing with dry or liquid fertilizer, do not exceed the recommended rate of 7 g/ha/application.

To make a 10% concentrated solution: dissolve 140 grams into 1 litre of water (20 oz/1 gallon).

CROP	APPLICATION RATE		TIMING / FREQUENCY
	10% Solution	Undiluted	
Nursery Stock (Containerized and field grown): deciduous and Evergreen trees, annuals, perennials, herbs, and vegetables	40 ml / 100 L water (1.25 oz / 20 gal)	N/A	Apply every 30 days or as needed commencing at transplant or 3-5 leaf stage.
Fruits and Berries: plums, apricots, cherry, peach, blueberries, blackberries, raspberries, strawberries	50 ml / ha (0.7 oz / acre)	7 g / ha (0.1 oz / acre)	3-4 applications commencing at 75% petal drop and repeating with 21-28 day intervals
Apples and Pears			3-4 applications commencing at 50% flowering and repeating with 21 day intervals
Fruiting vegetables: tomatoes, peppers, beans, peas, onions			First application prior to fruit set and repeat after 2 weeks
Lettuce, cauliflower, cabbage, Chinese cabbage			First application at 3-5 leaf stage and repeat after 2 weeks
Potatoes			First application at tuber initiation and repeat after 2 weeks
Cucurbits: pumpkins, squash, melons, cucumbers			First application at flowering and repeat after 21 days
Grapes: table and wine			Apply every 4-6 weeks commencing once plants begin to grow
Herbs: basil, chives, coriander, dill, parsley, sage			Apply every 4-6 weeks commencing once plants begin to grow
Field crops: barley, corn, oats, soybean, sugar beet, wheat			After plants reach the 3-5 leaf stage. Use with sufficient water to cover.
Grasses grown for seed, sod, pasture, forage, alfalfa			Apply when 2-4" tall. Use with sufficient water to cover.
Compost and Soil	40 ml / 100 L of soil/compost	N/A	Apply to soil or compost and mix thoroughly.
Hydroponics and Cannabis	N/A	1 g/200 L water	Apply to roots weekly through grow and bloom, beginning at transplant

WARNING
Keep out of reach of children and animals:
 Harmful if swallowed. Causes serious eye damage. Avoid breathing dust, spray or mist. Use only outdoors or in a well-ventilated location. Wear protective gloves, clothing, eye and NIOSH/OSHA approved respirator during product mixing and loading. Do not eat, drink or smoke when using this product. Wash thoroughly after handling. Avoid entry into water sources. Do not let product enter drains when rinsing or disposing of equipment wash water. Immediately contact a health care provider if exposed, concerned or feeling unwell. If swallowed, rinse mouth. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If inhaled, move person to fresh air. Store locked up in cool, dry, well-ventilated place out of direct sunlight. Dispose of contents and packaging in accordance with local, regional and federal requirements.

Lot Number:

Expiry Date:

25 KG